arXiv:1207.5388v1 [hep-th] 23 Jul 2012

Universal stress-tensor correlation functions of strongly coupled conformal fluids

Ram Brustein$^{(1)}$, A.J.M. Medved$^{(2)}$

(1) Department of Physics, Ben-Gurion University, Beer-Sheva 84105, Israel
(2) Department of Physics & Electronics, Rhodes University, Grahamstown 6140, South Africa

ramyb@bgu.ac.il, j.medved@ru.ac.za

Abstract

The gauge-gravity duality relates on-shell graviton multi-point functions in an anti-de Sitter black brane background to stress-tensor correlation functions of strongly coupled fluids. We show that, for conformal fluids whose gravitational dual is Einstein gravity, such correlation functions are universal in a particular kinematic region and depend only on the Mandelstam variable s. Thus, we extend the universality of the two-point function or, equivalently, the universality of the shear viscosity to entropy ratio to multi-point correlation functions. We also consider the leading corrections to Einstein gravity in string theory. The effect of these corrections is similarly universal, inducing a specific difference in the angular dependence of the stress-tensor correlation functions in the same kinematic region. We discuss schematically how multi-particle correlations in heavy-ion collisions can be used to test these findings and, thereby, probe the existence of a gravitational dual to the quark-gluon plasma and its universality class. We also propose a new way to measure the shear viscosity to entropy ratio using multi-point correlation functions.

1 Introduction

The gauge–gravity duality states that strongly coupled gauge theories in four dimensions have a dual description in terms of weakly coupled gravity theories in five-dimensional anti-de Sitter (5D AdS) space [1, 2]. In some sense, the gauge theory lives at the outer boundary of the AdS spacetime, and so this is a holographic correspondence [3].

If the bulk gravity theory is Einstein's, then all of its on-shell amplitudes depend on a single dimensionful parameter, the 5D Newton's constant G_5. Hence, appropriately chosen ratios of amplitudes do not depend on G_5. For example, the ratio of the shear viscosity to the entropy density, $\eta/s = 1/4\pi$ [4], is related to a ratio of two-point functions of gravitons [5] and thus independent of G_5. In string theory, the leading corrections to Einstein gravity are universal and depend on one additional dimensionful parameter, the string tension α'.

So far, the emphasis in theoretical and experimental studies has been on the two-point functions; the shear viscosity, conductivity, *etcetera*. From the experimental side, the prototypical example of a strongly coupled fluid is the quark–gluon plasma (QGP) [6]. The QGP is produced in heavy-ion collisions in the Large Hadron Collider (LHC) at CERN and the Relativistic Heavy Ion Collider (RHIC) at Brookhaven. Experiments on the QGP have already had significant success at testing the duality. In particular, the unusually low value of η/s was first predicted via the gauge–gravity duality [4] and then later substantiated [7, 8].

As multi-particle correlations in heavy-ion collisions have recently been measured [9, 10], the need for extracting information that can be utilized to probe the gravitational dual of the QGP has become pertinent. In the context of the gauge–gravity duality, the simplest field-theory objects are the connected correlation functions for the stress (energy-momentum) tensor. These are dual to graviton n-point functions [3].

We have recently calculated the one-particle irreducible (1PI) on-shell amplitudes of gravitons [11] and since extended the calculation to several connected n-point functions [12]. This work provides the basic ingredients to evaluate the connected stress-tensor correlators. We have focused on tensor-graviton amplitudes and on a certain kinematic region in which the amplitudes simplify significantly. The calculation was performed for Einstein gravity and for its leading-order corrected theory, Gauss–Bonnet gravity. We have identified the following universal properties:

1) All odd n-point functions of tensor gravitons vanish.

2) For Einstein gravity, the four- and higher-point functions all go as the Mandelstam center-of-mass variable s.

3) For the Gauss–Bonnet correction, the four-point function goes as s^2 and the higher-point functions go as $s(s+v)$, where v is another independent angle, interpreted as a "generalized Mandelstam variable".

2 The gravitational dual

We wish to explain briefly how on-shell graviton n-point functions are calculated. The details can be found in [11, 12].

The gravitational Lagrangian is expanded out in numbers of gravitons, where a graviton $h_{\mu\nu}$ represents a small perturbation of the metric from its background value, $g_{\mu\nu} \to g_{\mu\nu} + h_{\mu\nu}$. The $n^{\rm th}$ order of the expansion can be used to read off the graviton 1PI n-point function. Since we are eventually interested in the dual stress-tensor correlation functions, the bulk n-point functions are evaluated at some large but finite radius. We calculate amputated amplitudes in momentum space so the n-point functions are just functions of momentum.

In practice, one chooses the following ansatz for the gravitons: $h_{\mu\nu} = \phi(r)\exp\left[i\omega t - kz\right]$. Here, z indicates the direction of graviton propagation along the brane (or along any other spacetime slice at constant radius) and r is the usual AdS radial coordinate, ranging from $r = r_h$ at the horizon of the black brane to infinity at the AdS boundary. The remaining two spacelike directions will be denoted by x and y. For future reference, $a, b, \cdots = \{t, x, y, z\}$. We work in the radial gauge, for which $h_{rr} = h_{ar} = 0$ for any choice of a. This allows the gravitons to be separated into three distinct classes [13]: tensors, vectors and scalars. In the following, we will only be interested in the tensors h_{xy}.

The results are presented below for Einstein gravity in units for which $1/16\pi G_5$ is set to unity and $s = -k_1^{\mu} k_{2\mu}$ is the standard Mandelstam variable. The external gravitons are symmetrized.

We will be interested in the kinematic regime of "high momentum" in which the correlation functions of the stress-tensor simplify considerably. Also, the comparison between the leading Einstein result and possible corrections becomes simpler, as the number of derivatives in the interaction vertices is emphasized. In this regime, ω, k and s are large enough so as to dominate over all radial derivatives, whether with respect to the perturbative modes or the background. And so restricting to high momenta means that k^2, ω^2, $s \gg 1/L^2$, where L is the AdS curvature scale.

The simplest example is the two-point function, $\lim_{r\to\infty}\langle h_{xy}(k)h_{xy}(-k)\rangle_E = \left(\frac{L}{r}\right)^3 k^2$. And, in the high-momentum region, the higher-point functions for $2n = 4, 6, 8\ldots$ are given by

$$\lim_{r\to\infty}\langle (h_{xy})^{2n}\rangle^E_{1PI} \;=\; \frac{(2n-1)\Gamma\left[n+\frac{1}{2}\right]}{\sqrt{\pi}\Gamma[n-1]}\left(\frac{L}{r}\right)^{4n-1} s\ , \qquad (1)$$

with all the odd-point functions vanishing.

Rather than the above 1PI functions, we are really interested in connected functions of gravitons, as explained later. The connected functions can be calculated from the 1PI functions by evaluating tree-level Feynman diagrams, and these will have counterpart Witten diagrams on the boundary (see Fig. 1). However, we have found in [12] that, for Einstein gravity in the high-momentum regime, many of the connected functions are actually equal to the 1PI functions. We have checked that this persists up to the eight-point function and have strong indications that this phenomenon could persist for all the higher-point functions.

In addition to explicit calculations showing the equality of the 1PI and connected amplitudes, the following general argument further supports their equality. The generating function of the connected amplitudes is the Legendre transform of the generating function of the 1PI amplitudes. In general, there is no reason for these functions coincide. Einstein gravity in the high momentum regime is, however, special. General covariance and the universal appearance of two derivatives combine to severely restrict the form of either generating function [14, 11], and so they can be expected to be in agreement.

In any event, the difference between the connected and 1PI functions can only be in their numerical coefficients, while the angular (in)dependence of the amplitudes remain linear in s. So that,

$$\lim_{r\to\infty}\langle (h_{xy})^{2n}\rangle^{E}_{Con} \;=\; a_n \left(\frac{L}{r}\right)^{4n-1} \; s \;, \tag{2}$$

with $a_4 = 9/4$, $a_6 = 75/8$ and so on.

3 From bulk to boundary

We now want to translate the previous findings into statements about the dual strongly coupled fluid.

Since the stress (energy-momentum) tensor T_{ab} of the boundary field theory is dual to the bulk graviton h_{ab} [3], the connected on-shell graviton n-point functions can be used to calculate the (symmetrized) connected correlation function of the stress tensor, $\langle T_{a_1 b_1} T_{a_2 b_2} \dots T_{a_n b_n}\rangle_{Con}$.

We need to apply the standard rules of holographic renormalization [15, 16, 17] to the bulk n-point functions. First, evaluate the bulk n-point functions at some large but finite radius $r = r_0$. Second, multiply by an appropriate conformal factor Ω^q which is determined by the conformal dimensions of the operators. Third, subtract off any divergences. One then only retains the part that survives in the $r_0 \to \infty$ limit.

The conformal factor for the stress-tensor correlation function can be deduced from the asymptotic form of the metric. For an AdS brane geometry, $\lim\limits_{r\to\infty} ds^2 = -\frac{r^2}{L^2}\left[\eta_{ab}dx^a dx^b\right] + \frac{L^2}{r^2}dr^2$, which tells us that $\Omega = r/L$. The power is $q = \Delta - 3$, where $\Delta = \sum \Delta_i$ is a sum over conformal dimensions and the subtraction of 3 comes from the dimension of the metric determinant.

The operators are the gravitons for which $\Delta_h = 2$ and the derivatives for which $\Delta_\nabla = 1$.

Following the described procedure, we find that all powers of r and L are stripped away from our previous expressions. What is left is a quantity that is finite and well defined at the boundary. Hence, the first and third steps turn out to be of no direct consequence. Subleading contributions to the metric could be important in principle but, in our case, only show up in terms that are asymptotically vanishing.

The end result is that the field-theory correlation functions exhibit the same angular dependence as their bulk correspondents,

$$\langle (T_{xy})^{2n} \rangle^{E}_{Con} \;=\; a_{2n} s \;\; , \tag{3}$$

with all odd-point functions vanishing. The coefficients a_n are equal to their bulk counterparts from Eq. (2). We emphasize that our results are only valid in the high-momentum region. The full 3-point function for the stress tensor does not vanish, as must follow from conformal symmetry.

We remark that the 1PI functions for the stress-tensor are not very meaningful because of complications that arise in defining and separating the 1PI functions from the connected ones. (See [18] for a recent explanation.)

It is instructive to understand the kinematics of the high-momentum regime from the boundary point of view. The AdS scale L can be identified with the spatial extent of the fluid [19, 20], and so the meaning of the high-momentum region is that k^{-1}, ω^{-1} and $s^{-1/2}$ are smaller than the fluid size. Hence, one looks at correlation functions on scales less than the size of the fluid and does so by imposing an infrared cutoff on the momenta and frequencies.

To consistently apply the gauge–gravity duality in an AdS black hole (or brane) background requires that the horizon is larger than the AdS scale or, equivalently, hotter than the surrounding AdS space. In equation, $\pi T L \gg 1$. To understand the origin of this requirement, one considers geometries in which the boundary theory is defined on $S^3 \times R$. In this case, L is also the radius of the boundary sphere, and so an extra dimensionful parameter is introduced which breaks conformal invariance spontaneously in the same way that the temperature does. In this context, the ratio $\pi T L$ corresponds to the ratio of the size of the boundary fluid to the thermal wavelength.

Recall the high-momentum condition $kL \gg 1$ and, since we are also interested in the hydrodynamic limit, $k/\pi T \ll 1$. It follows that $\frac{1}{\pi T L} \ll \frac{kL}{\pi T L} \ll 1$ and the dimensionless parameter $\pi T L$, which measures how much hotter the BH is than AdS space, also determines the range of momenta that occupy the high-momentum regime.

4 Leading Correction

In the context of the gauge–gravity duality, one neglects $1/N$ corrections, so that the leading correction to on-shell amplitudes comes from four-derivative corrections to the Einstein Lagrangian. The physical content of these corrections depends on the coefficient of the Riemann-tensor-squared term, as the other four-derivative terms (Ricci-tensor-squared and Ricci-scalar-squared) can be eliminated with a suitable choice of field redefinitions [21, 22, 11]. Since our interest is in unitary theories whose equations of motion are at most second order in time derivatives, we use this freedom in field redefini-



Figure 1: Witten diagrams for stress-tensor correlation functions. The circle represents the AdS outer boundary, the dots depict interaction vertices, single lines correspond to undifferentiated gravitons and double lines correspond to differentiated gravitons. (a) A disconnected four-point function. (b) A 1PI four-point function at order ϵ^0. (c) A 1PI six-point function at order ϵ. (d) A 1PR connected six-point function at order ϵ^0. (e) A 1PR connected eight-point function at order ϵ. (f) A 1PR connected ten-point function at order ϵ^2.

tion to put the corrections in the Gauss–Bonnet form.

Let ϵ be a dimensionless parameter that measures the relative strength of the Gauss–Bonnet corrections compared to the leading Einstein result. Then $\epsilon \sim l_p^2/L^2$, where l_p is the 5D Planck length. The parameter ϵ can be determined by the coefficient of the Riemann-tensor-squared term in the Gauss–Bonnet combination. On the other hand, ϵ also appears in the ratio of the shear viscosity to the entropy density $\frac{\eta}{s} = \frac{1}{4\pi}[1-8\epsilon]$ [23, 24]. On dimensional grounds, a term in the bulk Lagrangian with $2(m+1)$ derivatives should go as ϵ^m. A sensible perturbative expansion then requires that $\epsilon \ll 1$.

Since Einstein and Gauss–Bonnet gravity both have a two-derivative linearized field equation, their real distinction comes from the non-linear interaction terms. Einstein gravity has only two-derivative vertices, a property that greatly restricts and simplifies its higher-point 1PI functions [14, 11]. In contrast, the four-point and ϵ-order higher-point functions of Gauss–Bonnet gravity have four derivatives and, at higher orders in ϵ, there can be many more. This distinction between different powers of ϵ is illustrated in Fig. 1.

We have found that the two extra derivatives in Gauss–Bonnet gravity lead to a more complicated angular dependence. However, the expansion to a given order in ϵ is still universal. For instance, to first order in ϵ, one obtains a specific angular dependence that is induced by the terms quartic in derivatives. Each additional order in ϵ brings in added complexity.

To first order in ϵ, we have found that

$$\begin{aligned}\lim_{r\to\infty}\langle (h_{xy})^{2n}\rangle^{GB}_{1PI} &= \lim_{r\to\infty}\langle (h_{xy})^{2n}\rangle^{E}_{1PI} \\ &+ \frac{2}{5}\epsilon\binom{2n}{4}\frac{\Gamma\left[n+\frac{3}{2}\right]}{\sqrt{\pi}\Gamma[n-1]}\left(\frac{L}{r}\right)^{4n+1} s(s+v)\ ,\end{aligned} \tag{4}$$

where $v = -k_1^{\mu}\sum_{j=5}^{2n} k_{j\mu}$ is a "generalized" Mandelstam variable which accounts for the additional number of gravitons that are involved in the scattering process. For the four-point function, one should set $v = 0$.

We have also found that, just like for Einstein gravity, all 1PR functions with a single internal line (*e.g.*, Fig. 1e) vanish up to first order in ϵ. Meaning that, at order ϵ, the connected functions are equal to the 1PI ones up to at least $2n = 6$. Then, because the ϵ-order theory is similarly constrained by general covariance and a strict number of derivatives, this agreement could very well persist for larger values of n.

Going from the bulk to the boundary, we have to holographically renormalize each of the two terms in Eq. (4) separately because of their different r dependence. Then

$$\langle (T_{xy})^{2n} \rangle^{GB}_{Con} \;=\; \langle (T_{xy})^{2n} \rangle^{E}_{Con} \;+\; \epsilon c_n s(s+v) \;, \tag{5}$$

such that $v = 0$ when $n = 2$ and the odd-number functions are vanishing. For example, $c_4 = 3/4$, $c_6 = 315/8$ and so on.

The relevance of the corrections to the interaction terms is determined by the value of $L^2 k^2 \epsilon \sim k^2 l_p^2$, whereas the consistency of treating string theory effectively as a theory of gravity requires that $k^2 l_p^2 < 1$. So that, if the momenta are “Planckian”, then the corrections become substantial. But, as the hydrodynamic approximation requires that $L^2 k^2 \epsilon \ll (\pi L T)^2 \epsilon$, it is really the value of $(\pi L T)^2 \epsilon$ which determines whether the higher-point functions will be substantially corrected. If $(\pi L T)^2 \epsilon \ll 1$, then the corrections will not appear in higher-point functions, only in the two-point function. This will be important when we discuss the possibility of experimentally determining the corrections.

5 Contact with experiment

The multi-point correlation functions of the stress (energy-momentum) tensor are the key to testing our ideas. Imagine a heavy-ion collision that creates a drop of QGP fluid. The drop expands and cools and then, after it freezes, a variety of decay products (mostly pions) stream outward. The energy, momentum and the velocity of the decay products can be measured (more easily for charged particles than for neutral ones), so that it is possible to measure the angular distribution of energy and momentum for a large number of particles emanating from the QGP fluid. The correlation functions of the stress-tensor components of the fluid well before freeze out can then be deduced with the standard methods, in a way similar to evaluating multiplicity correlation functions (*e.g.*, [25]).

Let us now consider the relevant components of the stress tensor $T_{xy} = T_{yx}$; these being dual to the tensor graviton modes. We are, assuming that the fluid flows in the z direction. Our objective is then to express these components in terms of simple observables.

For a conformal boundary theory, the stress-tensor can be expressed to leading order as (*e.g.* [26])

$$T_{ab} \;=\; \frac{\rho}{3}\left[4u_a u_b + \eta_{ab} - 4\pi\frac{\eta}{s}\frac{P^c_{\ a}P^d_{\ b}}{\pi T}\left(\partial_c u_d + \partial_d u_c - \frac{2}{3}\eta_{cd}\eta_{cd}\partial_e u^e\right)\right] , \qquad (6)$$

where u^a is the four-velocity of the fluid, $P^{ab} = \eta^{ab} + u^a u^b$ projects vectors onto directions perpendicular to u^a, ρ is the energy density, T is the temperature and η/s is the ratio of shear viscosity to entropy density. To arrive at this form, we have used the relations $p = \rho/3$ for the pressure and $s = (\rho+p)/T = (4/3)\rho/T$. Indices are raised or lowered with the Minkowski metric η_{ab}.

As the coordinates x and y are the spacelike directions perpendicular to u^a, we can calculate T_{xy} by disregarding the first two terms in (6) and then eliminating the diagonal components; the latter by contracting the stress-tensor with the "binormal vector" $\epsilon^a_{\ b} \equiv \{1 \text{ if } a \neq b \text{ ; } 0 \text{ if } a = b\ \}$. Then, defining $4\pi\frac{\eta}{s}\Phi \equiv T^{x}_{\ y}$ and a polarization tensor $\widehat{\epsilon}^{cd} \equiv \frac{1}{2}\epsilon^{ab}P^{c}_{\ a}P^{d}_{\ b}$, we have

$$\Phi \ = \ -\widehat{\epsilon}^{ab}\frac{\rho}{3}\left(\tilde{\partial}_a u_b + \tilde{\partial}_b u_a\right) \ , \tag{7}$$

where $\tilde{\partial}_a$ is a dimensionless spatial derivative, measured in units of the temperature $\tilde{\partial} = \frac{1}{\pi T}\partial$.

As the above equation makes clear, Φ can be determined by measuring the angular distribution of the energies, momenta and velocities of the collision products.

6 Scales, couplings and kinematics

As we are interested in applying the gauge–gravity duality to calculate stress-tensor correlation functions, it is necessary to insure that our analysis conforms with the duality's region of validity.

The holographic dictionary tells us that the number of colors N is related to the 't hooft coupling λ and the Yang-Mills coupling g_{YM} of the field theory as $\lambda = g_{YM}^2 N$. The first restriction comes from the large-N limit, for which $N \to \infty$, $g_{YM} \to 0$ such that λ is finite and large. This means that $1/N$ corrections are neglected while $1/\lambda$ corrections are considered small but finite. But, for the QGP, $\lambda \sim 10$ while $N = 3$ and there appears to be a conflict of interests. Nevertheless, as many features of the duality seem to be insensitive to this discrepancy, it is standard operating procedure to treat N as much larger than λ (see [27] for a recent discussion). We will adhere to this philosophy.

A further constraint comes from the hydrodynamic approximation, which is an implicit consequence of treating the QGP as a fluid. In this case, all momenta and frequencies are expected to be substantially smaller than the QGP temperature T (*cf*, the end of Section 3). However, let us adopt the optimistic viewpoint that the hydrodynamic regime can be loosened to $(k/\pi T)^2$, $(\omega/\pi T)^2 \lesssim 1$ and, within the context of our results, $s/(\pi T)^2 \lesssim 1$. For the QGP at the LHC, $T \sim 400$ MeV and $L \sim 7$ fm , so that $\pi T L \sim 15$ and the range in s is about 200. This means that the opportunity to distinguish different powers of s from data is open.

As discussed previously, the value of the dimensionless parameter $\epsilon(\pi LT)^2$ determines whether the corrections can be substantial in higher-point functions. For the QGP, using the above estimates and $\epsilon = 1/8(1-4\pi\eta/s)$, we find that $\epsilon(\pi LT)^2 \sim 25(1-4\pi\eta/s)$. Meaning that the higher-point functions are potentially sensitive to small corrections away from Einstein gravity.

The disconnected correlation functions (ignored so far) can be used for additional consistency checks when comparing our predictions with experiment. Consider, for example, the disconnected part of the four-point function (Fig. 1a): $\langle hhhh\rangle_{Dis} = \frac{4!}{(2!)^2}\langle hh\rangle^2$ or $\frac{\langle hhhh\rangle_{Dis}}{\langle hh\rangle} = 6\langle hh\rangle$. Clearly, one can use the value of the disconnected four-point function as an alternative means for ascertaining the two-point function and therefore η/s. This method works also for higher-point disconnected functions.

7 Conclusion

We have calculated the n-point correlation functions for the stress tensor of a strongly coupled conformal fluid via the gauge–gravity duality. We have explained schematically how to compare these results to statistical tests on the quark–gluon plasma at the LHC and what are the possible outcomes of such a comparison.

Given that the experimental tests are indeed carried out, there are three distinct possibilities: (i) The gravitational dual is shown, within errors, to be in the universality class of Einstein gravity, (ii) the gravitational dual is shown to be in the universality class of a corrected Einstein gravity or (iii) the correlation functions of QGP do not follow from either. Each of the three possibilities would provide very interesting information about the nature of the QGP and about the existence of a gravity dual. The first outcome is, of course, our expectation. The second would be a surprise and a wonderful opportunity to study "Planckian" physics with the QGP. The third outcome would indicate that, if the the quark–gluon plasma has a gravitational dual, it has to be in a more complicated universality class, with the implication that the low value of η/s is perhaps less significant in this context.

Acknowledgments

We thank Ofer Aharony, Misha Lublinsky, Sasha Milov, Kyriakos Papadodimas, Jurgen Schukraft, Sheer El-Showk, Raimond Snellings, Amos Yarom for helpful discussions. The research of RB was supported by the Israel Science Foundation grant no. 239/10. The research of AJMM received support from a Rhodes University Discretionary Grant.

References

[1] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” Adv. Theor. Math. Phys. **2**, 231 (1998) [Int. J. Theor. Phys. **38**, 1113 (1999)] [arXiv:hep-th/9711200].

[2] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, “Large N field theories, string theory and gravity,” Phys. Rept. **323**, 183 (2000) [arXiv:hep-th/9905111].

[3] E. Witten, “Anti-de Sitter space and holography,” Adv. Theor. Math. Phys. **2**, 253 (1998) [arXiv:hep-th/9802150];

[4] G. Policastro, D. T. Son and A. O. Starinets, “The shear viscosity of strongly coupled N = 4 supersymmetric Yang-Mills plasma,” Phys. Rev. Lett. **87**, 081601 (2001) [arXiv:hep-th/0104066].

[5] R. Brustein and A. J. M. Medved, “The ratio of shear viscosity to entropy density in generalized theories of gravity,” Phys. Rev. D **79**, 021901 (2009) [arXiv:0808.3498 [hep-th]].

[6] J. Casalderrey-Solana, H. Liu, D. Mateos, K. Rajagopal and U. A. Wiedemann, “Gauge/String Duality, Hot QCD and Heavy Ion Collisions,” arXiv:1101.0618 [hep-th].

[7] P. Kovtun, D. T. Son and A. O. Starinets, “Viscosity in strongly interacting quantum field theories from black hole physics,” Phys. Rev. Lett. **94**, 111601 (2005) [hep-th/0405231].

[8] T. Schafer and D. Teaney, "Nearly Perfect Fluidity: From Cold Atomic Gases to Hot Quark Gluon Plasmas," Rept. Prog. Phys. **72**, 126001 (2009) [arXiv:0904.3107 [hep-ph]].

[9] K. Aamodt *et al.* [ALICE Collaboration], "Higher harmonic anisotropic flow measurements of charged particles in Pb-Pb collisions at $\sqrt(s_{(NN)}) = 2.76$ TeV," Phys. Rev. Lett. **107**, 032301 (2011) [arXiv:1105.3865 [nucl-ex]].

[10] B. Abelev *et al.* [ALICE Collaboration], "Anisotropic flow of charged hadrons, pions and (anti-)protons measured at high transverse momentum in Pb-Pb collisions at $\sqrt(s_{(NN)}) = 2.76$ TeV," arXiv:1205.5761 [nucl-ex].

[11] R. Brustein and A. J. M. Medved, "Graviton n-point functions for UV-complete theories in Anti-de Sitter space," Phys. Rev. D **85**, 084028 (2012) [arXiv:1202.2221 [hep-th]].

[12] R. Brustein and A. J. M. Medved, article in preparation.

[13] G. Policastro, D. T. Son and A. O. Starinets, "From AdS/CFT correspondence to hydrodynamics," JHEP **0209**, 043 (2002) [arXiv:hep-th/0205052].

[14] D. M. Hofman, "Higher Derivative Gravity, Causality and Positivity of Energy in a UV complete QFT," Nucl. Phys. B **823**, 174 (2009) [arXiv:0907.1625 [hep-th]].

[15] J. de Boer, E. P. Verlinde and H. L. Verlinde, "On the holographic renormalization group," JHEP **0008**, 003 (2000) [arXiv:hep-th/9912012].

[16] K. Skenderis, “Lecture notes on holographic renormalization,” Class. Quant. Grav. **19**, 5849 (2002) [arXiv:hep-th/0209067].

[17] I. Papadimitriou and K. Skenderis, “AdS / CFT correspondence and geometry,” arXiv:hep-th/0404176.

[18] S. El-Showk and K. Papadodimas, “Emergent Spacetime and Holographic CFTs,” arXiv:1101.4163 [hep-th].

[19] E. Witten, “Anti-de Sitter space, thermal phase transition, and confinement in gauge theories,” Adv. Theor. Math. Phys. **2**, 505 (1998) [hep-th/9803131].

[20] S. S. Gubser, “Comparing the drag force on heavy quarks in N=4 super-Yang-Mills theory and QCD,” Phys. Rev. D **76**, 126003 (2007) [hep-th/0611272].

[21] G. ’t Hooft, “An algorithm for the poles at dimension 4 in the dimensional regularization procedure”, Nucl. Phys. B **62** 444 (1973).

[22] M .D. Pollock, “On the Field-Redefinition Theorem in Gravitational Theories”, Acta Phys. Pol. B **39** 1315 (2008).

[23] M. Brigante, H. Liu, R. C. Myers, S. Shenker and S. Yaida, “Viscosity Bound Violation in Higher Derivative Gravity,” Phys. Rev. D **77**, 126006 (2008) [arXiv:0712.0805 [hep-th]].

[24] Y. Kats and P. Petrov, “Effect of curvature squared corrections in AdS on the viscosity of the dual gauge theory,” JHEP **0901**, 044 (2009) [arXiv:0712.0743 [hep-th]].

[25] A. Bilandzic, R. Snellings and S. Voloshin, “Flow analysis with cumulants: Direct calculations,” Phys. Rev. C **83**, 044913 (2011) [arXiv:1010.0233 [nucl-ex]].

[26] D. T. Son and A. O. Starinets, “Viscosity, Black Holes, and Quantum Field Theory,” Ann. Rev. Nucl. Part. Sci. **57**, 95 (2007) [arXiv:0704.0240 [hep-th]]

[27] S. Aoki, K. Hashimoto and N. Iizuka, “Matrix Theory for Baryons: An Overview of Holographic QCD for Nuclear Physics”, arXiv:1203.5386 [hep-th].